


There is a threshold for Bisphenol A



- High-quality studies show that a clear threshold for BPA can be derived below which no adverse effects occur.
- BPA acts through receptor-mediated mechanisms, which have a threshold mode of action. There is no clear evidence of a non-monotonic dose-response (NMDR) relationship for the effects of BPA.


Together, the available data indicate that it is possible to identify a threshold for BPA toxicity.

A clear threshold for BPA in high-quality studies



High-quality studies conducted in rodents indicate a clear threshold for BPA toxicity above a no-observable-adverse-effect-level (NOAEL) range of 5–25 mg/kg bw/d BPA. [1,2,3,4,5] Reproductive and developmental effects, such as effects on fertility occur at high doses of BPA only and are accompanied by systemic toxicity. Below these thresholds no treatment related effects were observed. Developmental metabolic effects, neurotoxicity and immunotoxicity were not observed at any dose of BPA in these or other high-quality studies. [6,7,8,9,10,11,12]

BPA is active in 34% of ToxCast assays identified as having “receptor” or “receptor binding” as the assay target.



Receptor-mediated Assays with BPA Activity in ToxCast Database

- Androgen receptor in SRC non-receptor tyrosine kinases
- Constitutive androstane receptor
- Farnesoid X receptor and SRC non-receptor tyrosine kinases
- G protein-coupled receptor
- Glucocorticoid receptor
- Glutamate receptor
- Mineralocorticoid receptor
- Peroxisome proliferator-activated receptor
- Pregnane X receptor
- Progesterone receptor
- Receptor tyrosine kinases
- Sigma non-opioid intracellular receptor
- Sodium channel receptor

The ToxCast project is a high-throughput chemical screening program by the US EPA developed to identify *in vitro* assays and responses that are relevant to *in vivo* toxicity. Data from EU EPA ToxCast indicate that BPA is an estrogen receptor agonist and is active in multiple receptor-mediated assays [13]; such ligand-receptor mechanisms are indicative of a threshold mode of action. The European Safety Authority (EFSA) found no clear indication of a NMDR relationship for BPA across multiple endpoints, indicating that BPA does not cause adverse effects at low doses that are not caused at higher doses. EFSA stated that the available data do not suggest that NMDR relationships are relevant for the risk assessment of BPA. [14]

References for BPA Threshold Information Sheet



1. Tyl, RW; Myers, CB; Marr, MC; Thomas, BF; Keimowitz, AR; Brine, DR; Veselica, MM; Fail, PA; Chang, TY; Seely, JC; Joiner, RL; Butala, JH; Dimond, SS; Cagen, SZ; Shiotsuka, RN; Stropp, GD; Waechter, JM. 2002. "Three-generation reproductive toxicity study of dietary bisphenol A in CD Sprague-Dawley rats." *Toxicol. Sci.* 68:121-146.
2. Tyl, R; Myers, C; Marr, M; Sloan, C; Castillo, N; Veselica, M; Seely, J; Dimond, S; Van Miller, J; Shiotsuka, R; Beyer, D; Hentges, SG; Waechter, J. 2008. "Two-generation reproductive toxicity study of dietary bisphenol A (BPA) in CD-1® (Swiss) mice." *Toxicol. Sci.* 104(2):362-384.
3. Delclos, KB; Camacho, L; Lewis, SM; Vanlandingham, MM; Latendresse, JR; Olson, GR; Davis, KJ; Patton, RE; Gamboa da Costa, G; Woodling, KA; Bryant, MS; Chidambaram, M; Trbojevic, R; Juliar, BE; Felton, RP; Thorn, BT. 2014. "Toxicity evaluation of bisphenol A administered by gavage to Sprague-Dawley rats from gestation day 6 through postnatal day 90." *Toxicol. Sci.* 139(1):174-197.
4. Stump, DG; Beck, MJ; Radovsky, A; Garman, RH; Freshwater, LL; Sheets, LP; Marty, MS; Waechter, JM; Dimond, SS; Van Miller, JP; Shiotsuka, RN; Beyer, D; Chappelle, AH; Hentges, SG. 2010. "Developmental Neurotoxicity Study of Dietary Bisphenol A in Sprague Dawley Rats." *Toxicol. Sci.* 115(1):167-182.
5. Camacho, L; Lewis, SM; Vanlandingham, MM; Olson, GR; Davis, KJ; Patton, RE; Twaddle, NC; Doerge, DR; Churchwell, MI; Bryant, MS; McLellen, FM; Woodling, KA; Felton, RP; Maisha, MP; Juliar, BE; Gamboa da Costa, G; Delclos, KB. 2019. "A two-year toxicology study of bisphenol A (BPA) in Sprague-Dawley rats: CLARITY-BPA core study results." *Food Chem. Toxicol.* 132: 110728.
6. George, JD; Price, CJ; Tyl, RW; Marr, MC; Kimmel, C. 1985. "Teratologic evaluation of bisphenol A (CAS No. 80-05-7) administered to CD rats on gestational days 6 through 15." NTP Report.
7. Reel, JR; George, J; Lawton, AD; Myers, CB. 1985. "Bisphenol A: Reproduction and fertility assessment in CD-1 mice when administered in the feed." NTP Report.
8. Ema, M; Fujii, S; Furukawa, M; Kiguchi, M; Ikka, T; Harazono, A. 2001. "Rat two-generation reproductive toxicity study of bisphenol A." *Reprod. Toxicol.* 15:505-523.
9. Kobayashi, K; Ohtani, K; Kubota, H; Miyagawa, M. 2010. "Dietary exposure to low doses of bisphenol A: effects on reproduction and development in two generations of C57BL/6J mice." *Congenital Anomalies.* 50(3):159-170.
10. Kobayashi, K; Kubota, H; Ohtani, K; Hojo, R; Miyagawa, M. 2012. Lack of effects for dietary exposure of bisphenol A during in utero and lactational periods on reproductive development in rat offspring." *J. Toxicol. Sci.* 37(3):565-573.
11. Ferguson, SA; Law, CD; Abshire, JS. 2011. "Developmental treatment with bisphenol A or ethinyl estradiol causes few alterations on early preweaning measures." *Toxicol. Sci.* 124(1):149-160.
12. Ferguson, SA; Law, CD; Abshire, JS. 2012. "Developmental treatment with bisphenol A causes few alterations on measures of postweaning activity and learning." *Neurotoxicol. Teratol.* 34(6):598-606.
13. United States Environmental Protection Agency (US EPA). 2023. "US EPA ToxCast Data Release September 2023: Summary Files." 12p.
14. European Food Safety Authority (EFSA), Scientific Committee. 2021. "Opinion on the impact of non-monotonic dose responses on EFSA's human health risk assessments." *EFSA J.* 19(10):6877.